

# Flow cytometric determination of phytoplankton DNA in cultures and oceanic populations

Nicolas Boucher\*, Daniel Vaultot\*\*, Frédéric Partensky\*\*\*

CNRS et Université de Paris 6, Station Biologique BP 74, F-29682 Roscoff Cédex, France

**ABSTRACT:** A method to measure DNA and chlorophyll fluorescence simultaneously in single phytoplankton cells was developed and tested on cultured and natural populations. Samples were stored at  $-80^{\circ}\text{C}$  following fixation with 1% glutaraldehyde and freezing in liquid nitrogen, stained with DAPI (4',6-diamino-2-phenylindole), and analyzed by flow cytometry. In cultures, cell DAPI-DNA fluorescence measured by flow cytometry was related to cell DNA content measured by fluorometry with DAPI, over almost 4 orders of magnitude. In natural populations, the method easily discriminated photosynthetic cells from other living and non-living particles and permitted computation of the fraction of particulate DNA contained in photosynthetic picoplankton. In the northwestern Mediterranean Sea in summer, this fraction was higher at neritic than at pelagic stations, and at the latter was maximum at mid-depth. In the future, this method should permit (1) better estimates of biomass partitioning among the different trophic compartments, and (2) studies of cell cycling of oceanic plankton populations.

## INTRODUCTION

DNA is the most significant biological molecule on Earth because it encodes all living processes. Despite its small contribution to marine organic matter (ca 5% of 'particulate' and 0.1% of 'dissolved' organic carbon, respectively; Paul et al. 1985, 1988), it is an important biogeochemical marker which has been relatively little studied in aquatic environments (e.g. Holm-Hansen et al. 1968, Holm-Hansen 1969b, Pillai & Ganguly 1972, Falkowski & Owens 1982, Dortch et al. 1983, Paul et al. 1985, 1988, Winn & Karl 1986, Karl & Bailliff 1989).

In contrast to other cellular constituents, which vary widely with growth conditions, DNA content of a given cell type varies within a narrow range, typically 2-fold in extent for eukaryotes, with extremes reached at the beginning ( $G_1$  phase) and end ( $G_2$  phase) of the cell

cycle (Prescott 1976). Therefore, it has been hypothesized that particulate DNA concentrations could be used to provide reasonable estimates of living biomass (Holm-Hansen et al. 1968, Dortch et al. 1983). Indeed, cell DNA was found to be linearly related to cell carbon for a variety of phytoplankton species (Holm-Hansen 1969a). The failure of initial attempts (Holm-Hansen 1969b) to reconcile biomass estimates from DNA, on the one hand, and from particulate carbon, chlorophyll, or ATP, on the other hand, was attributed to the presence of an important fraction of 'detrital' DNA. This assertion has been challenged recently, at least in the case of coastal waters (Falkowski & Owens 1982, Dortch et al. 1983). Paul et al. (1985) found that particulate DNA was actually closely correlated with particulate organic carbon (POC) and bacterial counts in a wide range of environments. Winn & Karl (1986) noted, however, that only 10 to 25% of the particulate DNA appeared to be replicating actively in Pacific oligotrophic waters.

Some of the contradictions among these studies can be ascribed to the use of conversion factors (e.g. between carbon and DNA, carbon and ATP) that are assumed to apply universally to all compartments of the food web. This assumption might not hold, as recent measurements (Simon & Azam 1989) show that the DNA:carbon ratio is higher in bacteria than in phytoplankton and, for the former, decreases as cell size

\* Present address: Department of Biological Sciences, University of California at Santa Barbara, Santa Barbara, California 93106, USA

\*\* Addressee for reprint requests (Roscoff). On temporary leave at: Oceanography Department, University of Hawaii, 1000 Pope Road, Honolulu, Hawaii 96822, USA

\*\*\* On temporary leave at: Biological Oceanography Division, Bedford Institute of Oceanography, Dartmouth, Nova Scotia, Canada B2Y 4A2

increases. Therefore, in order to obtain better biomass estimates from DNA and to solve the riddle of 'detrital' DNA, it appears necessary to separate the contribution of the different communities to particulate DNA. One way to achieve this result is to perform filter-fractionation; Paul et al. (1985) have shown that in the Gulf of Mexico, between 70 and 99 % of the particulate DNA is contained in the 0.2 to 1  $\mu\text{m}$  fraction, suggesting that bacteria dominate in particulate DNA, at least in pelagic waters. However, the complexity of the microbial loop and the interpenetration between the size-classes of autotrophs and heterotrophs (as evidenced by the recent discovery of bacteria-sized primary producers, Chisholm et al. 1988) singularly complicate interpretation of such measurements.

A potential solution is to measure DNA in single cells. This may be easily done by flow cytometry, which also discriminates autotrophs from other cell types in natural populations, using the red chlorophyll fluorescence of the former (Olson et al. 1985), and is very sensitive since it has been used recently for bacterial DNA determination in aquatic samples (Robertson & Button 1989).

In the present paper, we describe a flow cytometric method to measure quantitatively DNA in natural phytoplankton populations, which is calibrated with bulk fluorometric DNA determinations performed on a set of phytoplankton strains. The method is applied to estimate the fraction of particulate DNA contained in photosynthetic picoplankton in the northwestern Mediterranean Sea in summer.

## MATERIAL AND METHODS

**Cultured species.** Eleven non-axenic phytoplankton strains belonging to 6 algal classes and representing a wide range of algal cell sizes (Table 1) were grown in either K (Keller et al. 1987) or f/2 (Guillard & Ryther

1962) medium. Batch cultures were maintained at 20°C in 500 ml Erlenmeyer flasks under 100  $\mu\text{E m}^{-2} \text{s}^{-1}$  of fluorescent light (Sylvania F58 WI-154ST) supplied on a 12 h light:12 h dark cycle. Cultures were sampled in late exponential phase to maximize yield and minimize bacterial contamination.

**Field samples.** Four stations were sampled with Niskin bottles at 4 to 6 levels depending on bottom depth during the EROS-2000 cruise of the Italian RV 'Bannock' in the northwestern Mediterranean Sea between 1 July and 15 July 1989.

**Cell densities and size.** For cultures, replicate ( $n > 4$ ) measurements were made on 0.5 ml samples using a Coulter Counter (model ZM, Coulter, Hialeah, Florida) equipped with a 100  $\mu\text{m}$  diameter orifice and coupled to a logarithm range expander and a C1000 Channelyser. Since *Synechococcus* spp. are too small to be sized accurately with the 100  $\mu\text{m}$  orifice, their abundance was determined by flow cytometry (see below) on replicate 100  $\mu\text{l}$  samples ( $n > 4$ ).

**Particulate DNA.** DNA was quantified using the method developed by Brunk et al. (1979) and modified by Falkowsky & Owens (1982). In brief, samples of cultured (20 to 50 ml) or natural (350 ml) populations were gently (< 100 mm Hg) filtered onto Whatman GF/F filters washed with a 3% NaCl, 10 mM EDTA solution. When the analysis could not be performed immediately (e.g. natural samples), filters were frozen in liquid nitrogen and stored at -80°C. Filters were homogenized on ice in 2 ml of a solution (A) containing 100 mM NaCl (5.85 g l<sup>-1</sup>), 10 mM EDTA (3.35 g l<sup>-1</sup>) and 100 mM Tris (12.1 g l<sup>-1</sup>) at pH 7. The slurry was centrifuged 5 min at 1500 rpm. The DNA-containing supernatant was kept at 4°C until analysis, which was routinely performed within 3 h.

Stock solutions of DAPI (4',6-diamino-2-phenylindole, mol. wt = 279; Sigma, St. Louis, Missouri, D1388) were prepared in solution A at a final concentration of 100 ng ml<sup>-1</sup> (0.36  $\mu\text{M}$ ). Standard DNA stock solution

Table 1. Cultured algal species used in this study

Class	Species	Strain	Laboratory of origin
Cyanophyceae	<i>Synechococcus</i> sp.	ROS04	Roscoff (France)
	<i>Synechococcus</i> sp.	WH7803	CCMP-Bigelow (USA)
Cryptophyceae	<i>Cryptomonas maculata</i>		Plymouth (UK)
Dinophyceae	<i>Amphidinium carteri</i>	Amphi	CCMP-Bigelow (USA)
	<i>Gymnodinium</i> cf. <i>nagasakiense</i>	Tinduff	Roscoff (France)
	<i>Gymnodinium nagasakiense</i>	Buzen-85-2	Kitasato (Japan)
Prymnesiophyceae	<i>Isochrysis galbana</i>	Iso	L'Houmeau (France)
	<i>Hymenomonas carterae</i>	CoccoII	CCMP-Bigelow (USA)
	<i>Phaeocystis pouchetii</i>		Texel (Netherlands)
Bacillariophyceae	<i>Thalassiosira weissflogii</i>	Actin	CCMP-Bigelow (USA)
Chlorophyceae	<i>Dunaliella primolecta</i>		Plymouth (UK)

(final concentration  $20 \mu\text{g ml}^{-1}$ ) was obtained by dissolving calf thymus DNA (Sigma Type I, D1501) at room temperature in solution A (pH = 10) and adjusting to pH 7 with Tris-HCl.

Fluorescence measurements (excitation = 350 nm; emission = 460 nm) were performed on a JY3D spectrofluorometer (Jobin et Yvon, France) equipped with a 150 W xenon light source, and DNA concentration was computed as detailed in Falkowski & Owens (1982).

**Flow cytometric DNA determination.** Unconcentrated samples (1 ml) were fixed with 1% glutaraldehyde (electron microscopy grade; Merck, Darmstadt, Germany), incubated for 10 min at room temperature in the dark, frozen in liquid nitrogen (Vaulot et al. 1989), and then stored at  $-80^\circ\text{C}$ . Prior to analysis, samples were thawed at room temperature, stained with DAPI to a final concentration of  $1 \mu\text{g ml}^{-1}$  ( $3.6 \mu\text{M}$ , cultures) or  $2 \mu\text{g ml}^{-1}$  ( $7.2 \mu\text{M}$ , natural samples) and incubated for 90 min at room temperature in the dark. Calf thymocyte nuclei (Fluorotrol RF; Ortho Diagnostic System, Inc., Westwood, Massachusetts) and  $2 \mu\text{m}$  low-intensity Nile Red beads (Pandex, Mundelein, Illinois) were used as external (cultures) or internal (natural samples) standards.

For each sample 100 to 300  $\mu\text{l}$  were analyzed on an Epics 541 (Coulter) flow cytometer fitted with a 6 W argon laser (Model 90-6; Coherent, Palo Alto, California) and a Biosense flow cell. Laser emission was set at 351/364 nm (UV mode) and 100 mW (cultures) or 200 mW (natural samples). Scattered light (a sizing parameter) was measured at forward (FALS) and right (RALS) angles. RALS was collected after reflection on a 400 nm long-pass (LP) dichroic filter (MTO, France). Emitted fluorescence was filtered through a 415 nm LP filter (Coulter) and split with a 590 nm short-pass dichroic filter (Coulter) to separate blue DAPI-DNA fluorescence, collected through a 485 nm band-pass filter (Glen Spectra, Stanmore, UK), and red chlorophyll fluorescence, collected through a 690 nm LP filter (MTO). All 4 parameters (FALS, RALS, blue DAPI-DNA and red chlorophyll fluorescences) were recorded in peak (culture) or integral (for improved sensitivity in natural samples) mode on 3-decade logarithmic range (256 channels). Natural samples were recorded in list mode. Data were processed on an IBM-PC-compatible microcomputer using custom-designed software, CytoPC. Briefly, each picoplankton population (prochlorophytes, *Synechococcus* spp., eukaryotes; see 'Results') was identified by its flow cytometric signature on light-scatter vs chlorophyll cytograms and discriminated from the others by a rectangular gate. Its DNA histogram was obtained by playing back the list mode data and keeping only the cells that fell within the gate. From this histogram, the average DAPI-DNA fluorescence of the population was

computed, all cell cycle stages being included, and was normalized to the fluorescence of the calf thymocyte nuclei.

## RESULTS AND DISCUSSION

### Phytoplankton DNA staining for flow cytometry

The first step in developing a DNA staining method for natural phytoplankton populations was to find an appropriate fixative. Its functions are (1) to preserve the cells so as to allow delayed analysis, (2) to preserve chlorophyll red fluorescence in order to differentiate photosynthetic cells from other particles, and (3) to make the cell membrane permeable so as to facilitate stain penetration. In cell cycle studies, alcohol preservation (methanol or ethanol) is by far the most widespread technique, but it cannot be used for natural populations since it extracts photosynthetic pigments (Olson et al. 1983). A method combining 1% glutaraldehyde fixation and liquid nitrogen freezing has been recently shown to preserve phytoplankton adequately for flow cytometric analyses (Vaulot et al. 1989). Preliminary tests showed that cells preserved with this method were amenable to DNA staining.

The second step was to choose a fluorescent DNA stain among the dozen available (Shapiro 1985), meeting several criteria. First, the fluorescence emission wavelengths of the stain and of the photosynthetic pigments must be distinct, so that the 2 fluorescences can be separated optically. A significant overlap of the emission bands results in a strong correlation of the 2 emitted fluorescences (cross-talk). This phenomenon prevents the use of red-emitting stains (e.g. propidium iodide) for chlorophyll-containing cells, and that of orange-emitting ones (e.g. chromomycin A3) for phycoerythrin-containing cells (cyanobacteria, Cryptophyceae). Second, the stain must bind stoichiometrically to DNA over the wide range of cell DNA content encountered in phytoplankton (typically from 0.1 to 100 pg cell $^{-1}$ ; Holm-Hansen 1969a). Stoichiometry is also essential for cell cycle studies when cells in  $G_1$  have to be distinguished from those in  $G_2 + M$  (Olson et al. 1983). Third, the fluorescence yield of the stain must be large enough to allow for analysis of the small picoplankton populations which dominate in most natural samples. With these criteria in mind, DAPI seemed to be a good choice. It is specific for DNA and does not require treatment with RNase, in contrast to propidium iodide, which binds to both RNA and DNA (Shapiro 1985). Its emission peak (around 460 nm) clearly falls out of both phycoerythrin and chlorophyll emission bands. It has a high fluorescence yield and as such has been recently used to stain the small bacteria encoun-




Fig. 1. *Thalassiosira weissflogii*. Flow cytometric analysis of the diatom fixed with 1% glutaraldehyde, frozen in liquid nitrogen, and stained with  $1 \mu\text{g ml}^{-1}$  of DAPI. Contoured cytogram: distribution of red chlorophyll fluorescence vs DAPI-DNA fluorescence, both on 3-decade logarithmic scales. The 2 populations correspond to the  $G_1$  and  $G_2 + M$  phases of the cell cycle. Insert: DNA distribution on a linear scale

tered in aquatic systems (Robertson & Button 1989). One drawback of DAPI is that its affinity for DNA depends on the base pair composition, since it binds specifically to AT pairs (Lin et al. 1977).

We settled therefore for the following procedure: 1% glutaraldehyde fixation (final pH = 6.6), freezing in liquid nitrogen, and staining with  $1 \mu\text{g ml}^{-1}$  DAPI. This protocol was found to stain all species effectively. Fluorescence was well above background, as the blue autofluorescence of cells in a non-stained sample never exceeded 5% of that of stained cells. Saturation was reached within 1 h. There was no detectable interference between chlorophyll and DAPI fluorescences, as demonstrated by the absence of correlation between the 2 parameters in cytograms (Fig. 1). The coefficient of variation (CV) of the  $G_1$  peak ranged from 10 to 36% (average = 19.5%,  $n = 11$ ), with the worst results obtained for *Synechococcus* sp. Although we were able to resolve the  $G_1$  and  $G_2$  peaks in most cases (Fig. 1), the CVs obtained with this method were larger than those obtained with methanol fixation. For example, we obtained a CV of 13.4% for *Thalassiosira weissflogii*, while Vaultot et al. (1986) reported a CV of 6.5% with the latter method. Large CVs are probably a consequence of imperfect stain penetration rather than absorbance of DAPI fluorescence by chlorophyll, as suggested by Olson et al. (1983); the latter phenomenon would induce a negative correlation between chlorophyll and DAPI fluorescences, never observed in our case.

Absence of fixation or of cell freezing at low temperature resulted in very poor stain penetration, as observed previously (Olson et al. 1983). Fixation with either buffered glutaraldehyde (pH = 7.5) or paraformaldehyde often resulted in lower DAPI-DNA fluorescence yields. The improvement at low pH is probably due to partial removal of histones (Giangarè et al. 1989).

In most species, stain penetration was facilitated by treatment of the cell membrane with detergent (0.1% Triton X100; Traganos et al. 1977). This increased DAPI-DNA fluorescence up to 30% and separated the  $G_1$  and  $G_2$  peaks better. Detergent treatment induced extrusion of cytoplasmic material and cell aggregation in some cases (e.g. *Dunaliella primolecta*) and was therefore not applicable to all species.

### Standardization and calibration

In order to relate single-cell DAPI-fluorescence to cell DNA content, flow cytometric fluorescence measurements must first be standardized in order to eliminate day-to-day variations in instrument optimization. This is usually achieved with fluorescent beads. In the present case, reference cell nuclei are more indicated since they will also reflect day-to-day variations in staining conditions. We employed calf thymocyte nuclei, which are widely used and commercially available. It would also have been possible to use a given phytoplankton species to standardize all the others. When few samples are analyzed (e.g. cultures), reference cells can be used as external standards, provided they are stained under exactly the same conditions as the cells (i.e. in the present case in seawater with 1% glutaraldehyde). When a large number of samples are

Table 2. Cultured species: diameters, measured with a Coulter Counter, and DNA content, measured with DAPI by fluorometry. DNA content is the average over all cell cycle phases and not that of  $G_1$  phase cells

Species	Diameter ( $\mu\text{m}$ )	DNA content ( $\text{pg cell}^{-1}$ )
<i>Synechococcus</i> ROS04	1.5 <sup>a</sup>	0.027
<i>Synechococcus</i> WH7803	1.5 <sup>a</sup>	0.022
<i>Cryptomonas maculata</i>	7.8	1.5
<i>Amphidinium carteri</i>	6.7	6.5
<i>Gymnodinium</i> cf. <i>nagasakiense</i>	19.4	71
<i>Gymnodinium nagasakiense</i>	21.1	123
<i>Isochrysis galbana</i>	4.0	0.42
<i>Hymenomonas carterae</i>	8.6	1.72
<i>Phaeocystis pouchetii</i>	5.1	0.43
<i>Thalassiosira weissflogii</i>	15.0	4.70
<i>Dunaliella primolecta</i>	5.9	0.72
<sup>a</sup> Estimated		




Fig. 2. Cell DAPI-DNA fluorescence measured by flow cytometry (normalized to calf thymocyte nuclei; per thymocyte) vs cell DNA content ( $\text{pg cell}^{-1}$ ), measured by fluorometry for different phytoplankton strains (Tables 1 & 2). (A) Linear scale ( $n = 9$ ). (B) Logarithmic scale ( $n = 11$ ). Regression lines are given in the text by Eqs. (1) and (2), respectively. Open symbols correspond to dinoflagellates

to be run (natural populations), it is simpler to use them as internal standards, i.e. mixed with the sample.

In order to calibrate standardized flow cytometric measurements, we compared them to cellular DNA concentrations ( $\text{DNA}_c$  in  $\text{pg cell}^{-1}$ ) established by fluorometry using DAPI (Falkowski & Owens 1982) for 11 strains of phytoplankton (Table 2). On a linear scale, we found a very good correlation between the 2 measurements in the range 0 to  $7 \text{ pg cell}^{-1}$  for  $\text{DNA}_c$ , excluding the 2 *Gymnodinium* species, which have a very high DNA content (Fig. 2A):

$$\text{DNA}_c = 17.4 \times \text{DNA}_f + 0.06 \quad (r^2 = 1.00, n = 9) \quad (1)$$

where  $\text{DNA}_f$  is blue DAPI-DNA fluorescence standardized to that of calf thymocyte nuclei.

Since  $\text{DNA}_c$  spanned almost 4 orders of magnitude for the set of strains considered (Table 2), it appeared appropriate to plot these data on logarithmic scales (Fig. 2B):

$$\text{DNA}_c = 13.9 \times \text{DNA}_f^{0.92} \quad (r^2 = 0.99, n = 11) \quad (2)$$

The good correlations obtained in both cases indicate that, provided the same stain is used (DAPI), the flow cytometric method, for which DNA is maintained packaged inside the cells, and the fluorometric method, for which the DNA is extracted, yield very comparable results. The slight non-linearity of Eq. (2) (exponent = 0.92 instead of 1) indicates that larger cells are more readily stained than smaller cells. It is noteworthy that dinoflagellates, although their DNA structure is very different from that of the other eukaryotes (Rizzo 1987), did not depart from the regression line. If  $\text{DNA}_f$  is set equal to 1,  $\text{DNA}_c$  is equal to 17.5 (Eq. 1) or 13.9 (Eq. 2)  $\text{pg cell}^{-1}$ , which should represent the DNA content of the thymocytes. It is higher, however, than the actual

value ( $7.3 \text{ pg cell}^{-1}$ ; Tiersch et al. 1989) and reflects the fact that bare thymocyte nuclei are more easily stained than glutaraldehyde-fixed phytoplankton cells.

As mentioned earlier, DAPI fluorescence is a function of both absolute DNA content and base pair composition. Therefore, if instead of measuring cell DNA fluorometrically with DAPI we had measured it with a stain which is not base-sensitive, such as 3,5-diaminobenzoic acid (DABA; Holm-Hansen 1969a) or ethidium bromide, we would have observed a lower correlation with the flow cytometric measurements, since base composition varies widely in phytoplankton




Fig. 3. Cell DNA content measured by fluorometry vs cell volume for 11 phytoplankton strains (Tables 1 & 2). Regression line is given in the text by Eq. (3) and does not include prokaryotes (*Synechococcus* sp., 2 points at lower left) nor dinoflagellates (open symbols). The other line corresponds to the relation established by Shuter et al. (1983)

even within a single species. For example, in *Chlorella sorokiniana* G + C content varies between 61.5 and 75.2 % (Huss et al. 1986).

Mean cell volume ( $V$ , in  $\mu\text{m}^3$ ) was found to be well correlated with cell DNA content, if prokaryotes and Dinophyceae were excluded (Fig. 3):

$$V = 164 \times \text{DNA}_c^{1.46} \quad (r^2 = 0.97, n = 6) \quad (3)$$

Such a relation is qualitatively similar to that established by Shuter et al. (1983) for a wide range of unicellular eukaryotes and suggests that DNA might be a good estimator of phytoplankton biomass. Part of the discrepancy between our data (Fig. 3) and those of Shuter et al. (1983) could come from the heterogeneity of their data set, where cell volume and cell DNA have often been obtained from different investigators. Prokaryotes were not included in Eq. (3), because they exhibit a volume-to-DNA relation very different from that of eukaryotes (Shuter et al. 1983). The latter authors also found that dinoflagellates fitted well to the eukaryotic relation, while our data indicate that the volume:DNA ratio was 10 times smaller for dinoflagellates than for other species. Although more measurements are clearly needed before drawing definitive conclusions, our data are actually consistent with the fact that this algal class is well known for its large number of chromosomes and its high proportion of structural DNA, which compensates for the low abundance of nuclear histone-like proteins (Rizzo 1987). Moreover, DNA content may vary widely among dinoflagellate species of similar size. This is the case with the morphologically similar taxa *Gymnodinium nagasakiense*, widespread in Japanese waters, and *Gymnodinium* cf. *nagasakiense*, common in European seas, which exhibit a 2-fold discrepancy in DNA contents (Table 2) despite equivalent numbers of chromosomes (Partensky et al. 1988).

#### Application to natural populations

Several problems may arise when a staining method, tested on cultures, is applied to natural populations. First, cells can be lost due to fixation and preservation. In the present case, however, the fixation method has been extensively tested on natural and cultured populations and appears to induce minimal cell loss, except for some fragile phytoplankton classes such as cryptophytes and dinoflagellates (Vaulot et al. 1989). In contrast, alcohol fixation, which requires several rounds of centrifugation, results in over 90 % cell loss (D. Vaulot unpubl.). Second, natural samples contain a whole array of particles that might be stained non-specifically by the fluorochrome and interfere with detection of the cells of interest. Third, species that are




Fig. 4. Stns BA8, BA17, BA18 and BA20, sampled in the northwestern Mediterranean Sea in July 1989

maintained in culture are not fully representative of those that thrive in the oceans (Giovannoni et al. 1990), and the response of the most fragile species can only be tested with natural samples.

To examine these potential problems, the method was tested on natural picoplankton populations from samples taken in summer at 4 stations located on an offshore transect in the northwestern Mediterranean Sea (Fig. 4). In a typical sample, 3 picoplankton populations of increasing cell size and decreasing abundance were usually present: prochlorophytes (Chisholm et al. 1988), *Synechococcus* spp. cyanobacteria, and picoeukaryotes (Fig. 5A). The identity of each group was established by parallel flow cytometric analysis of the sample using visible light (488 nm) and, for the prochlorophytes, by spectrofluorometric pigment analysis (Vaulot et al. 1990). The 3 populations were stained correctly by our procedure (Fig. 5B). Non-living particles and bacteria proved not to be a major problem: although they interfered with prochlorophytes in ungated cytograms (Fig. 5B), they could be easily removed by gating the picoplankton populations on the scatter vs chlorophyll cytograms (Fig. 5C). As in cultured populations, there was no interference between DAPI-DNA and chlorophyll fluorescences, as visualized by the absence of correlation between the 2 parameters in the corresponding cytograms (Fig. 5C). Eukaryotes were usually composed of 2 sub-populations. One gave sharp DNA distributions with CVs as low as 9 % (Fig. 5C and middle eukaryotic peak in Fig.




Fig. 5. Phytoplankton sampled from Stn BA18, 55 m depth, in the northwestern Mediterranean Sea. (A) Cytogram of right angle light scatter (sizing parameter) vs chlorophyll fluorescence (log units, 3 decades). Three cell populations are visible: prochlorophytes ( $46\,800\text{ cell ml}^{-1}$ ), *Synechococcus* spp. cyanobacteria ( $16\,600\text{ cell ml}^{-1}$ ), and picoeukaryotes ( $2\,370\text{ cell ml}^{-1}$ ). The population on the lower right-hand side with large scatter and low fluorescence is composed of detrital particles. Calf thymocyte nuclei and  $2\ \mu\text{m}$  beads have been added as internal standards. The dashed rectangle represents a gating region containing only photosynthetic picoplankton. (B) Cytogram of DAPI-DNA fluorescence vs chlorophyll fluorescence (log units, 3 decades). Note interference between the particles and the prochlorophytes. The apparent high red fluorescence of the thymocyte nuclei is a consequence of their very high blue DAPI fluorescence (it would fall around 10 000, out of scale, on the x-axis), which leaks into the red channel. (C) Same as (B) but only for populations contained within the gate drawn in (A). Note that gating removed the interfering particles as well as the  $2\ \mu\text{m}$  beads and the thymocyte nuclei. (D). DNA histograms for the different cell populations. Note that vertical axis is in log units. Picoplankton DNA concentration was estimated, using Eq. (2), to be equal to  $2.3\ \mu\text{g l}^{-1}$  in this sample

5D) and probably was constituting a single taxonomic entity; the other had a broad DNA distribution and was likely a mixture of several cell types. *Synechococcus* spp. had DNA distributions with non-distinct peaks and CVs on the order of 25 %, similar to those observed for cultures (Fig. 5D). The very small prochlorophytes ( $0.4$  to  $0.8\ \mu\text{m}$  in diameter) produced a very good DNA

signal with sharp peaks having CVs on the order of 10 % (Fig. 5D). Their DAPI-DNA fluorescence was actually more intense than for *Synechococcus* spp., despite the larger size of the latter (Fig. 5A). Although prochlorophytes could have a larger DNA content (and likely a larger DNA:C ratio) or be richer in AT base pairs, this was probably a consequence of easier stain

penetration in prochlorophytes: *Synechococcus* spp. cyanobacteria are notorious for their very tough membrane (Bricaud et al. 1988), and while Triton X100 treatment increases stain penetration in *Synechococcus* sp., it lyses prochlorophytes (data not shown).

Using Eq. (2) to convert DAPI-DNA fluorescence to DNA content for each population, photosynthetic picoplankton DNA was estimated and compared to total particulate DNA (Table 3). It must be stressed that these estimates are not fully representative of the total phytoplankton community since they do not include the larger plankton (typically  $> 2 \mu\text{m}$ ), which was probably in significant concentration near the coast (in surface samples, cells  $> 1 \mu\text{m}$  accounted for 85 % of the chlorophyll *a* at Stn BA8 and for 62 % at Stn BA18; J. Neveux pers. comm.). The relative contribution of the 3 picoplanktonic groups was variable: prochlorophytes dominated at the 3 stations closer to shore, and *Synechococcus* at the outermost station BA20 (Table 3). In some cases, we obtained unreasonably high estimates of picoplankton DNA (e.g. Stn BA18, 55 m), which might be due to an overestimate of prochlorophyte DNA because of their good stainability (see above).

When all samples were pooled together, no correlation was observed between total particulate DNA and picoplankton DNA ( $r^2 = 0.036$ ,  $n = 28$ ,  $P = 0.331$ ), in agreement with previous studies that failed to find any significant relation between particulate DNA and chlorophyll (Holm-Hansen 1969b, Paul et al. 1985). This points out the highly variable contribution of phytoplankton to particulate organic matter. A clearer pattern emerged if neritic and pelagic stations were contrasted (Table 3). At neritic stations (BA8, BA17) picoplankton accounted for a higher proportion of particulate DNA, in accordance with the observations of Falkowski & Owens (1982) and Paul et al. (1985). At

pelagic stations (BA18, BA20) picoplankton contribution was in general lower, especially in the upper euphotic zone and at depth, while it peaked at the picoplankton maximum. This corroborates previous reports (Paul et al. 1985, Winn & Karl 1986, Fuhrman et al. 1989, Cho & Azam 1990) asserting that, in oligotrophic waters, phytoplankton is a small fraction of the total living biomass.

## CONCLUSION

The method described in this paper should permit quantitative determination of DNA in single phytoplankton cells. It is much more informative than total particulate DNA measurements, since it allows assessment of how DNA is partitioned among the different compartments of the food web. In the present study, we restricted ourselves to picoplankton. However, flow cytometry and DAPI staining have already helped characterize bacterial communities (Robertson & Button 1989) and can certainly be applied to larger phytoplankton ( $> 2 \mu\text{m}$ ) and microzooplankton, using instruments designed to handle larger sample volumes (Dubelaar et al. 1989).

One source of error in cell DNA estimates by flow cytometry may arise from differences in stain penetration among species. Although this was not observed in cultures, as evidenced by the very good fit obtained for Eq. (2), we suspect that it occurred for field populations, in particular for prochlorophytes, which had a much higher DAPI fluorescence than expected on the basis of their size. Therefore, calibration should be performed using cultured species as closely related as possible to the field populations of interest.

A supplemental source of error comes from the base specificity of DAPI. Consider a cell population whose

Table 3. Mean  $\pm$  SD of total particulate DNA (measured by fluorometry) and DNA in 3 photosynthetic picoplankton populations (measured by flow cytometry and computed as detailed in text) in different depth strata in the northwestern Mediterranean Sea in July 1989 (Fig. 4). For neritic stations (BA8, BA17), where there was no definite pattern in the vertical distribution of picoplankton DNA, all depths were pooled together. For pelagic stations (BA18, BA20), 3 layers were distinguished: upper euphotic zone, picoplankton maximum, and below the picoplankton maximum

Stn	Depth stratum (m)	Particulate DNA ( $\mu\text{g l}^{-1}$ )	Prochlorophyte DNA ( $\mu\text{g l}^{-1}$ )	<i>Synechococcus</i> spp. DNA ( $\mu\text{g l}^{-1}$ )	Picoeukaryotes DNA ( $\mu\text{g l}^{-1}$ )	Picoplankton DNA (%)	n
BA8 <sup>a</sup>	5–70	3.2 $\pm$ 2.5	1.00 $\pm$ 0.35	0.30 $\pm$ 0.19	0.53 $\pm$ 0.48	68 $\pm$ 18	3
BA17	8–70	2.4 $\pm$ 1.4	0.88 $\pm$ 0.17	0.19 $\pm$ 0.06	0.28 $\pm$ 0.06	74 $\pm$ 44	4
BA18	0–15	2.6 $\pm$ 1.0	0.13 $\pm$ 0.06	0.10 $\pm$ 0.01	0.24 $\pm$ 0.07	20 $\pm$ 7	3
	55	2.0	1.70	0.31	0.26	110	1
	145–295	13.7 $\pm$ 2.7	0.01 $\pm$ 0.00	0.01 $\pm$ 0.01	0.02 $\pm$ 0.01	0 $\pm$ 0	2
BA20	5–30	6.5 $\pm$ 3.6	0.28 $\pm$ 0.12	0.28 $\pm$ 0.13	0.30 $\pm$ 0.04	14 $\pm$ 3	3
	50	8.3	0.42	1.74	0.45	31	1
	75–100	8.7 $\pm$ 4.3	0.00 $\pm$ 0.00	0.05 $\pm$ 0.05	0.06 $\pm$ 0.04	1 $\pm$ 0	2

<sup>a</sup>Value from 45 m not taken into account

contribution to particulate DNA is  $D_1$  ( $\mu\text{g l}^{-1}$ ) and whose AT base composition is  $f_1$  (a number between 0 and 1).  $D_2$  and  $f_2$  correspond to the rest of the living biomass. Assuming that  $D_2 \gg D_1$ , i.e. that the cell population is only a small fraction of the total biomass, and that the fluorescence of DAPI is proportional to the product  $f_1 D_1$  (although this might be an oversimplified assumption, as it has been recently shown that DAPI binds to a consensus of 4 consecutive AT base pairs; Portugal & Waring 1988), then the flow cytometric estimate of the fraction of DNA contained in the cell population would be equal to  $f_1 D_1 / f_2 D_2$ , while the actual value is  $D_1 / D_2$ . The maximum range of variation for the AT fraction among all living organisms lies between 0.20 and 0.80 (Stanier et al. 1970). If we take the median of this range for  $f_2$  (0.5), the flow cytometric estimate would fall between 0.4 and 1.6 times the actual value.

Despite these limitations, the linear relation registered between cell volume and DNA (Eq. 3) indicates that the latter could be used to estimate biomass, once rigorous calibration has been performed. Future studies should compare such estimates with carbon estimates.

The use of DNA in flow cytometric analyses of natural picoplankton populations may also help assess their heterogeneity in conjunction with other cell properties such as pigment fluorescence characteristics (Olson et al. 1988), antibody-binding (Shapiro et al. 1989), or oligonucleotide-tagging (Amann et al. 1990). For example, prochlorophytes and *Synechococcus* spp. are better separated on DNA vs chlorophyll cytograms (Fig. 5C) than on scatter vs chlorophyll (Fig. 5A). Increased separation might also be obtained by using, in conjunction with the AT-binding DAPI, a GC-binding dye such as chromomycin A3 (Van Dilla et al. 1983).

Another benefit of measuring single-cell DNA in natural populations is to gain information on their cell cycle status. Numerous laboratory studies have investigated how the cell cycle of marine and freshwater phytoplankton is controlled by environmental factors such as light, temperature, or nutrients (cf. reviews in Puiseux Dao 1981, Donnan et al. 1985). However, these controls have never been demonstrated in the field. The very good staining ability of the prochlorophytes (see above) enabled us to actually analyze their cell cycle in natural samples and to demonstrate its control by nitrogen in situ (Vaulot & Partensky unpubl.). For other cell types such as *Synechococcus* sp. (Fig. 5), our procedure will have to be improved (e.g. by detergent addition) to yield a better definition of the cell cycle phases. Flow cytometric measurements of DNA in natural phytoplankton populations will facilitate the application of a recently developed method to derive growth rates of synchronized populations from time series of DNA distributions (Carpenter & Chang 1988),

providing a datum much sought after to resolve the long-standing controversy about the exact magnitude of phytoplankton growth rates in oligotrophic waters (Goldman et al. 1979).

*Acknowledgements.* Help from Claude Courties and Jean-Louis Birrien is kindly acknowledged, as well as constructive comments from 2 referees. The EPICS 541 was funded by the Institut National des Sciences de l'Univers (CNRS). Financial support for field work was provided by the Commission of the European Communities 4th Environment R&D Programme under contract EV4V-011-F (EROS-2000 Project).

#### LITERATURE CITED

- Amann, R. I., Binder, B. J., Olson, R. J., Chisholm, S. W., Devereux, R., Stahl, D. A. (1990). Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for analyzing mixed microbial populations. *Appl. Environ. Microbiol.* 56: 1919-1925
- Bricaud, A., Bédhomme, A.-L., Morel, A. (1988). Optical properties of diverse phytoplanktonic species: experimental results and theoretical interpretation. *J. Plankton Res.* 10: 851-873
- Brunk, C. F., Jones, K. C., James, T. W. (1979). Assay for nanogram quantities of DNA in cellular homogenates. *Analyt. Biochem.* 92: 497-500
- Carpenter, E. J., Chang, J. (1988). Species-specific phytoplankton growth rates via diel DNA synthesis cycles. I. Concept of the method. *Mar. Ecol. Prog. Ser.* 43: 105-111
- Chisholm, S. W., Olson, R. J., Zettler, E. R., Waterbury, J., Goericke, R., Welschmeyer, N. (1988). A novel free-living prochlorophyte occurs at high cell concentrations in the oceanic euphotic zone. *Nature, Lond.* 334: 340-343
- Cho, B. C., Azam, F. (1990). Biogeochemical significance of bacterial biomass in the ocean's euphotic zone. *Mar. Ecol. Prog. Ser.* 63: 253-259
- Donnan, L., Carvill, E. P., Gilliland, T. J., John, P. C. L. (1985). The cell cycles of *Chlamydomonas* and *Chlorella*. *New Phytol.* 99: 1-40
- Dortch, Q., Roberts, T. L., Clayton, J. R. Jr, Ahmed, S. I. (1983). RNA/DNA ratios and DNA concentrations as indicators of growth rate and biomass in planktonic marine organisms. *Mar. Ecol. Prog. Ser.* 13: 61-71
- Dubelaar, G. B., Groenewegen, A. C., Stokdijk, W., van den Engh, G. J., Visser, J. W. (1989). Optical plankton analyser: a flow cytometer for plankton analysis. II. Specifications. *Cytometry (N.Y.)* 10: 529-539
- Falkowski, P. G., Owens, T. C. (1982). A technique for estimating phytoplankton division rates by using a DNA-binding fluorescent dye. *Limnol. Oceanogr.* 27: 776-782
- Fuhrman, J. A., Sleeter, T. D., Carlson, C. A., Proctor, L. M. (1989). Dominance of bacterial biomass in the Sargasso Sea and its ecological implications. *Mar. Ecol. Prog. Ser.* 57: 207-217
- Giangarè, M. C., Prosperi, E., Pedraly-Noy, G., Bottiroli, G. (1989). Flow cytometric evaluation of DNA stainability with propidium iodide after histone H1 extraction. *Cytometry (N.Y.)* 10: 726-730
- Giovannoni, S. J., Britschgi, T. B., Moyer, C. L., Field, K. G. (1990). Genetic diversity in Sargasso Sea bacterioplankton. *Nature, Lond.* 345: 60-63
- Goldman, J. C., McCarthy, J. J., Peavey, D. W. (1979). Growth rate influence on the chemical composition of phytoplankton in oceanic waters. *Nature, Lond.* 279: 210-215

- Guillard, R. R., Ryther, J. H. (1962). Studies of marine diatoms. I. *Cyclotella nana* Husdelt and *Detonula confervacea* Gran. Can. J. Microbiol. 8: 229–239
- Holm-Hansen, O. (1969a). Amount of DNA and organic carbon in single cells. Science 63: 87–88
- Holm-Hansen, O. (1969b). Determination of microbial biomass in ocean profiles. Limnol. Oceanogr. 14: 740–747
- Holm-Hansen, O., Sutcliffe, W. H. Jr, Sharp, J. (1968). Measurements of deoxyribonucleic acid in the ocean and its ecological significance. Limnol. Oceanogr. 13: 507–514
- Huss, V. A. R., Dörr, R., Grossmann, U., Kessler, E. (1986). Deoxyribonucleic acid reassociation in the taxonomy of the genus *Chlorella*. I. *Chlorella sorokiniana*. Archs Microbiol. 145: 329–333
- Karl, D. M., Bailliff, M. D. (1989). The measurement and distribution of dissolved nucleic acids in aquatic environments. Limnol. Oceanogr. 34: 543–558
- Keller, M. D., Selvin, R. C., Claus, W., Guillard, R. R. L. (1987). Media for the culture of oceanic ultraphytoplankton. J. Phycol. 23: 633–638
- Lin, M. S., Comings, D. E., Alfi, O. S. (1977). Optical studies of the interaction of 4'-6-diamino-2-phenylindole with DNA and metaphase chromosomes. Chromosoma 60: 15–25
- Olson, R. J., Chisholm, S. W., Zettler, E. R., Armbrust, E. V. (1988). Analysis of *Synechococcus* pigment types in the sea using single and dual beam flow cytometry. Deep Sea Res. 35: 425–440
- Olson, R. J., Frankel, S. L., Chisholm, S. W., Shapiro, H. M. (1983). An inexpensive flow cytometer for the analysis of fluorescence signals in phytoplankton: chlorophyll and DNA distributions. J. exp. mar. Biol. Ecol. 68: 129–144
- Olson, R. J., Vault, D., Chisholm, S. W. (1985). Marine phytoplankton distributions measured using shipboard flow cytometry. Deep Sea Res. 32: 1273–1280
- Partensky, F., Vault, D., Couté, A., Sournia, A. (1988). Morphological and nuclear analysis of the bloom forming dinoflagellates *Gyrodinium cf. aureolum* and *Gymnodinium nagasakiense* (Dinophyceae). J. Phycol. 24: 408–415
- Paul, J. H., DeFlaun, M. F., Jeffrey, W. H., David, A. W. (1988). Seasonal and diel variability in dissolved DNA and in microbial biomass and activity in a subtropical estuary. Appl. environ. Microbiol. 54: 718–727
- Paul, J. H., Jeffrey, W. H., DeFlaun, M. (1985). Particulate DNA in subtropical oceanic and estuarine planktonic environments. Mar. Biol. 90: 95–101
- Pillai, T. N. V., Ganguly, A. K. (1972). Nucleic acids in the dissolved constituents of sea-water. J. mar. biol. Ass. India 14: 384–390
- Portugal, J., Waring, M. J. (1988). Assignment of DNA binding sites to 4',6-diamidino-2-phenylindole and bisbenzimidazole (Hoechst 33258). A comparative footprinting study. Biochim. biophys. Acta 949: 158–168
- Prescott, D. M. (1976). Reproduction in eukaryotic cells. Academic Press, New York
- Puiseux-Dao, S. (1981). Cell-cycle events in unicellular algae. In: Platt, T. (ed.) Can. Bull. Fish. Aquat. Sci. 210: 130–149
- Rizzo, P. J. (1987). Biochemistry of the dinoflagellate nucleus. In: Taylor, F. J. R. (ed.) Botanical Monographs, Vol. 21, The biology of dinoflagellates. Blackwell Scientific Publications, Oxford, p. 143–173
- Robertson, B. R., Button, D. K. (1989). Characterizing aquatic bacteria according to population, cell size, and apparent DNA content by flow cytometry. Cytometry (N.Y.) 10: 70–76
- Shapiro, H. M. (1985). Practical flow cytometry. Alan R. Liss, Inc., New York
- Shapiro, L. P., Campbell, L., Haugen, E. M. (1989). Immunochemical recognition of phytoplankton species. Mar. Ecol. Prog. Ser. 57: 219–224
- Shuter, B. J., Thomas, J. E., Taylor, W. D., Zimmerman, A. M. (1983). Phenotypic correlates of genomic DNA content in unicellular eukaryotes and other cells. Am. Nat. 122: 26–44
- Simon, M., Azam, F. (1989). Protein content and protein synthesis rates of planktonic bacteria. Mar. Ecol. Prog. Ser. 51: 201–213
- Stanier, R. Y., Doudoroff, M., Adelberg, E. A. (1970). The microbial world, 3rd edn. Prentice Hall, Inc., Englewood Cliffs, New Jersey
- Tiersch, T. R., Chandler, R. W., Wachtel, S. S., Elias, S. (1989). Reference standards for flow cytometry and application in comparative studies of nuclear DNA content. Cytometry (N.Y.) 10: 706–710
- Traganos, F., Darzynkiewicz, Z., Sharpless, T., Melamed, M. R. (1977). Simultaneous staining of ribonucleic and deoxyribonucleic acids in unfixed cells using acridine orange in a flow cytofluorometric system. J. Histochem. Cytochem. 25: 46–56
- Van Dilla, M. A., Langlois, R. G., Pinkel, D., Yajko, D., Hadley, W. K. (1983). Bacterial characterization by flow cytometry. Science 220: 620–622
- Vault, D., Courties, C., Partensky, F. (1989). A simple method to preserve oceanic phytoplankton for flow cytometric analyses. Cytometry (N.Y.) 10: 629–635
- Vault, D., Olson, R. J., Chisholm, S. W. (1986). Light and dark control of the cell cycle in two phytoplankton species. Expl. Cell Res. 167: 38–52
- Vault, D., Partensky, F., Neveux, J., Mantoura, R. F. C., Llewellyn, C. (1990). Winter presence of prochlorophytes in surface waters of the northwestern Mediterranean Sea. Limnol. Oceanogr. 35: 1156–1164
- Winn, C. D., Karl, D. M. (1986). Diel nucleic acid synthesis and particulate DNA concentrations: conflicts with division rate estimates by DNA accumulation. Limnol. Oceanogr. 31: 637–645.

This article was presented by M. Levandowsky, New York, New York, USA

Manuscript first received: July 30, 1990  
Revised version accepted: January 24, 1991